



---

# Lymphocyte T-Cell Immunomodulator

## TREATMENT INSTRUCTIONS

**ADMINISTRATION:** 1 ml dosage (1µg/ml) by subcutaneous injection.

**RECOMMENDED TREATMENT SCHEDULE:** Upon diagnosis of FeLV or FIV infection.

- **Month 1:** Weekly injections
- **Month 2:** Every other week
- **Month 3 +:** Every 4 - 6 weeks or as necessary

Patient responses may vary, and some patients may require more frequent treatments.

**PATIENT MONITORING:**

**Clinical Signs:**

The most important monitoring method is clinical improvement. Each patient should be monitored for clinical signs associated with their infection(s). IMULAN has established a standard clinical evaluation and scoring form to monitor patient progress. This form is available upon request.

**Blood Counts:**

When LTCI is used as an aid in the treatment of FeLV or FIV disease, monitoring of blood cell parameters, especially lymphocyte and red cell counts is recommended to evaluate the subject's progress. CBCs may be done every 4 weeks for patients that have lymphopenia and/or anemia.

**Other Methods:**

Other methods are available that can detect shifts in the types of lymphocytes, such as CD4+ or CD8 cells. Please contact IMULAN for assistance on additional testing at (877) 892-9595.

---

Phone Toll Free: 877-892-9595

Fax Toll Free: 866-603-9526



Visit: [www.IMULAN.com](http://www.IMULAN.com)

email: [info@IMULAN.com](mailto:info@IMULAN.com)